


# NORTHWEST HIV CARE REGION




**Population Counts, Northwest HIV Care Region, 2017**


County	White	Black/African American	Hispanic	Asian/Pacific Islander	American Indian/Alaskan Native	Two or More Races/Other	Total
Andrew County	16,617 94.7%	138 0.8%	430 2.4%	101 0.6%	45 0.3%	224 1.3%	17,555
Atchison County	5,069 96.1%	22 0.4%	71 1.3%	41 0.8%	14 0.3%	58 1.1%	5,275
Buchanan County	73,810 82.9%	5,128 5.8%	6,044 6.8%	1,587 1.8%	372 0.4%	2,124 2.4%	89,065
Caldwell County	8,606 94.6%	60 0.7%	192 2.1%	31 0.3%	50 0.5%	161 1.8%	9,100
Carroll County	8,327 94.7%	159 1.8%	139 1.6%	28 0.3%	24 0.3%	119 1.4%	8,796
Daviess County	7,998 95.7%	49 0.6%	137 1.6%	20 0.2%	35 0.4%	122 1.5%	8,361
DeKalb County	10,585 84.1%	1,479 11.7%	315 2.5%	56 0.4%	51 0.4%	102 0.8%	12,588
Gentry County	6,374 95.6%	40 0.6%	118 1.8%	31 0.5%	16 0.2%	86 1.3%	6,665
Grundy County	9,389 94.4%	89 0.9%	217 2.2%	75 0.8%	44 0.4%	135 1.4%	9,949
Harrison County	8,117 95.2%	45 0.5%	190 2.2%	39 0.5%	33 0.4%	100 1.2%	8,524
Holt County	4,231 95.9%	14 0.3%	58 1.3%	16 0.4%	49 1.1%	45 1.0%	4,413
Livingston County	13,972 92.1%	538 3.5%	304 2.0%	84 0.6%	68 0.4%	207 1.4%	15,173
Mercer County	3,478 94.6%	13 0.4%	96 2.6%	30 0.8%	18 0.5%	43 1.2%	3,678
Nodaway County	20,770 92.4%	643 2.9%	394 1.8%	373 1.7%	57 0.3%	235 1.0%	22,472
Worth County	1,987 96.6%	19 0.9%	29 1.4%	5 0.2%	6 0.3%	11 0.5%	2,057
<b>Region Total</b>	<b>199,330 89.1%</b>	<b>8,436 3.8%</b>	<b>8,734 3.9%</b>	<b>2,517 1.1%</b>	<b>882 0.4%</b>	<b>3,772 1.7%</b>	<b>223,671</b>

This page was intentionally left blank.

**Figure 1. HIV disease cases (living and deceased), by current HIV vs. stage 3 (AIDS) status, Northwest HIV Care Region, 1982-2018**



**Figure 2. Living and new HIV disease cases and deaths, by year\*, Northwest HIV Care Region, 2009-2018**




\*Living HIV disease cases represent the number of individuals living with HIV disease at the end of the year. New HIV disease cases represent the number of individuals newly diagnosed in the year. HIV disease deaths represent the number of individuals that died in the year.

From 1982 to 2018, there have been 239 HIV disease cases diagnosed in the Northwest HIV Care Region and reported to DHSS (Figure 1). Of the cumulative cases reported, 54.8% were still presumed to be living with HIV disease at the end of 2018. Among those living with HIV disease, 60 were classified as HIV cases at the end of 2018 and 71 were classified as stage 3 (AIDS) cases.


At the end of 2018, there were 131 persons living with HIV disease whose most recent diagnosis occurred in the Northwest HIV Care Region (Figure 2). The number of people living with HIV disease generally increased over time. There were eight new HIV disease diagnoses in 2018. The number of new diagnoses remained generally stable since 2009 with increases observed from 2012 to 2013 and from 2015 through 2017. The number of deaths among persons with HIV disease remained stable.

**Figure 3. HIV disease cases, by current status\* and year of diagnosis\*\*, Northwest HIV Care Region, 2009-2018**



\*HIV case vs. stage 3 (AIDS) case  
 \*\*Cases are indicated by year of initial diagnosis reported to DHSS (i.e., the year in which the first diagnosis of the person, whether as an HIV case or a stage 3 (AIDS) case, was documented by DHSS).  
 \*\*\*These cases were either: 1) initially reported as HIV cases and then later reclassified as stage 3 (AIDS) cases because they subsequently met the stage 3 (AIDS) case definition; or 2) initially reported as stage 3 (AIDS) cases.  
 \*\*\*\*These cases were initially reported as HIV cases and have remained HIV cases. They have not met the case definition for stage 3 (AIDS) as of December 31, 2018.


**Figure 4. Reported HIV disease cases, by current status\* and year of diagnosis\*\*, Northwest HIV Care Region, 2009-2018**



\*HIV case vs. stage 3 (AIDS) case.  
 \*\*Cases are indicated by year of initial diagnosis reported to DHSS (i.e., the year in which the first diagnosis of the person, whether as an HIV case or a stage 3 (AIDS) case, was documented by DHSS).  
 \*\*\*These cases were either: 1) initially reported as HIV cases and then later reclassified as stage 3 (AIDS) cases because they subsequently met the stage 3 (AIDS) case definition; or 2) initially reported as stage 3 (AIDS) cases.  
 \*\*\*\*These cases were initially reported as HIV cases and have remained HIV cases. They have not met the case definition for stage 3 (AIDS) as of December 31, 2018.

The number of new diagnoses remained generally stable since 2009 with the exception of increases observed from 2012 to 2013 and from 2015 through 2017 (Figures 3 and 4). There were decreases observed from 2009 to 2010, 2013 to 2015, and 2017 to 2018. Differences in the number of persons sub-classified as stage 3 (AIDS) cases each year are due to the progression of the disease over time.

**Figure 5. Persons diagnosed with HIV disease by current vital status\* and year of diagnosis\*\*, Northwest HIV Care Region, 2009-2018**



\*Vital status on December 31, 2018.

\*\*Cases are indicated by year of initial diagnosis reported to DHSS (i.e., the year in which the first diagnosis of the person, whether as an HIV case or a stage 3 (AIDS) case, was documented by DHSS).

Of the thirteen persons diagnosed with HIV disease in 2009, three (23.1%) were deceased by the end of 2018 (Figure 5). Of the eight individuals first diagnosed in 2018, all were alive by the end of 2018. Due to low overall numbers of new cases for each year, trends in HIV disease deaths are not easy to interpret.

**Table 1. Living<sup>†</sup> HIV, stage 3 (AIDS), and HIV disease cases, by sex, by race/ethnicity, by race/ethnicity and sex, and by current age, Northwest HIV Care Region, 2018**

	HIV*			Stage 3 (AIDS)**			HIV Disease***		
	Cases	%	Rate****	Cases	%	Rate****	Cases	%	Rate****
<b>Sex</b>									
Male	48	80.0%	42.7	49	69.0%	43.5	97	74.0%	86.2
Female	12	20.0%	10.8	22	31.0%	19.8	34	26.0%	30.6
<b>Total</b>	<b>60</b>	<b>100.0%</b>	<b>26.8</b>	<b>71</b>	<b>100.0%</b>	<b>31.7</b>	<b>131</b>	<b>100.0%</b>	<b>58.6</b>
<b>Race/Ethnicity</b>									
White	47	78.3%	23.6	50	70.4%	25.1	97	74.0%	48.7
Black/African American	9	15.0%	106.7	18	25.4%	213.4	27	20.6%	320.1
Hispanic	4	6.7%	45.8	3	4.2%	34.3	7	5.3%	80.1
Asian/Pacific Islander	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
American Indian/Alaskan Native	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
Two or More Races/Unknown	0	0.0%	--	0	0.0%	--	0	0.0%	--
<b>Total</b>	<b>60</b>	<b>100.0%</b>	<b>26.8</b>	<b>71</b>	<b>100.0%</b>	<b>31.7</b>	<b>131</b>	<b>100.0%</b>	<b>58.6</b>
<b>Race/Ethnicity-Males</b>									
White Male	41	85.4%	41.6	39	79.6%	39.6	80	82.5%	81.2
Black/African American Male	5	10.4%	89.8	7	14.3%	125.8	12	12.4%	215.6
Hispanic Male	2	4.2%	42.0	3	6.1%	63.0	5	5.2%	105.0
Asian/Pacific Islander Male	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
American Indian/Alaskan Native Male	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
Two or More Races/Unknown Male	0	0.0%	--	0	0.0%	--	0	0.0%	--
<b>Total</b>	<b>48</b>	<b>100.0%</b>	<b>42.7</b>	<b>49</b>	<b>100.0%</b>	<b>43.5</b>	<b>97</b>	<b>100.0%</b>	<b>86.2</b>
<b>Race/Ethnicity-Females</b>									
White Female	6	50.0%	6.0	11	50.0%	10.9	17	50.0%	16.9
Black/African American Female	4	33.3%	139.4	11	50.0%	383.3	15	44.1%	522.6
Hispanic Female	2	16.7%	50.4	0	0.0%	0.0	2	5.9%	50.4
Asian/Pacific Islander Female	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
American Indian/Alaskan Native Female	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
Two or More Races/Unknown Female	0	0.0%	--	0	0.0%	--	0	0.0%	--
<b>Total</b>	<b>12</b>	<b>100.0%</b>	<b>10.8</b>	<b>22</b>	<b>100.0%</b>	<b>19.8</b>	<b>34</b>	<b>100.0%</b>	<b>30.6</b>
<b>Current Age<sup>‡</sup></b>									
<2	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
2-12	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
13-18	0	0.0%	0.0	1	1.4%	5.9	1	0.8%	5.9
19-24	5	8.3%	24.9	2	2.8%	10.0	7	5.3%	34.9
25-44	31	51.7%	56.8	15	21.1%	27.5	46	35.1%	84.4
45-64	17	28.3%	29.7	45	63.4%	78.5	62	47.3%	108.2
65+	7	11.7%	17.6	8	11.3%	20.1	15	11.5%	37.7
<b>Total</b>	<b>60</b>	<b>100.0%</b>	<b>26.8</b>	<b>71</b>	<b>100.0%</b>	<b>31.7</b>	<b>131</b>	<b>100.0%</b>	<b>58.6</b>

<sup>†</sup>Includes persons diagnosed with HIV disease in the Northwest HIV Care Region who are currently living, regardless of current residence.

\*Cases which remained HIV cases at the end of 2018.

\*\*Cases classified as stage 3 (AIDS) by December 31, 2018.

\*\*\*The sum of HIV cases and stage 3 (AIDS) cases.

\*\*\*\*Per 100,000 population based on 2017 DHSS estimates.

<sup>‡</sup>Based on age as of December 31, 2018.

Note: Percentages may not total 100% due to rounding.

**Table 2. Diagnosed HIV, stage 3 (AIDS), and HIV disease cases, by sex, by race/ethnicity, by race/ethnicity and sex, and by current age, Northwest HIV Care Region, 2018**

	HIV*			Stage 3 (AIDS)**			HIV Disease***		
	Cases	%	Rate****	Cases	%	Rate****	Cases	%	Rate****
<b>Sex</b>									
Male	0	0.0%	0.0	4	100.0%	3.6	4	50.0%	3.6
Female	4	100.0%	3.6	0	0.0%	0.0	4	50.0%	3.6
<b>Total</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>8</b>	<b>100.0%</b>	<b>3.6</b>
<b>Race/Ethnicity</b>									
White	1	25.0%	0.5	3	75.0%	1.5	4	50.0%	2.0
Black/African American	1	25.0%	11.9	0	0.0%	0.0	1	12.5%	11.9
Hispanic	2	50.0%	22.9	1	25.0%	11.4	3	37.5%	34.3
Asian/Pacific Islander	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
American Indian/Alaskan Native	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
Two or More Races/Unknown	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	--
<b>Total</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>8</b>	<b>100.0%</b>	<b>3.6</b>
<b>Race/Ethnicity-Males</b>									
White Male	0	--	0.0	3	75.0%	3.0	3	75.0%	3.0
Black/African American Male	0	--	0.0	0	0.0%	0.0	0	0.0%	0.0
Hispanic Male	0	--	0.0	1	25.0%	21.0	1	25.0%	21.0
Asian/Pacific Islander Male	0	--	0.0	0	0.0%	0.0	0	0.0%	0.0
American Indian/Alaskan Native Male	0	--	0.0	0	0.0%	0.0	0	0.0%	0.0
Two or More Races/Unknown Male	0	--	0.0	0	0.0%	0.0	0	0.0%	0.0
<b>Total</b>	<b>0</b>	<b>--</b>	<b>0.0</b>	<b>4</b>	<b>100.0%</b>	<b>3.6</b>	<b>4</b>	<b>100.0%</b>	<b>3.6</b>
<b>Race/Ethnicity-Females</b>									
White Female	1	25.0%	1.0	0	--	0.0	1	25.0%	1.0
Black/African American Female	1	25.0%	34.8	0	--	0.0	1	25.0%	34.8
Hispanic Female	2	50.0%	50.4	0	--	0.0	2	50.0%	50.4
Asian/Pacific Islander Female	0	0.0%	0.0	0	--	0.0	0	0.0%	0.0
American Indian/Alaskan Native Female	0	0.0%	0.0	0	--	0.0	0	0.0%	0.0
Two or More Races/Unknown Female	0	0.0%	0.0	0	--	0.0	0	0.0%	--
<b>Total</b>	<b>4</b>	<b>100.0%</b>	<b>3.6</b>	<b>0</b>	<b>--</b>	<b>0.0</b>	<b>4</b>	<b>100.0%</b>	<b>3.6</b>
<b>Current Age<sup>†</sup></b>									
<2	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
2-12	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
13-18	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
19-24	0	0.0%	0.0	1	25.0%	5.0	1	12.5%	5.0
25-44	4	100.0%	7.3	1	25.0%	1.8	5	62.5%	9.2
45-64	0	0.0%	0.0	2	50.0%	3.5	2	25.0%	3.5
65+	0	0.0%	0.0	0	0.0%	0.0	0	0.0%	0.0
<b>Total</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>8</b>	<b>100.0%</b>	<b>3.6</b>

\*HIV cases diagnosed during 2018 which remained HIV cases at the end of the year.

\*\*Stage 3 (AIDS) cases initially diagnosed in 2018.

\*\*\*The sum of newly diagnosed HIV cases and newly diagnosed stage 3 (AIDS) cases. Does not include cases diagnosed prior to 2018 with HIV, which progressed to stage 3 (AIDS) in 2018.

\*\*\*\*Per 100,000 population based on 2017 DHSS estimates.

†Based on age as of December 31, 2018.


Note: Percentages may not total 100% due to rounding.

**Epi Profiles Summary: Northwest HIV Care Region**

Of the 131 persons living with HIV disease at the end of 2018, 74.0% were males (Table 1). The rate of those living with HIV disease among males was 2.8 times as high as the rate among females. Although whites represented the largest proportion of living HIV disease cases (74.0%), the rate of those living with HIV disease among blacks/African Americans was 6.6 times as high as the rate among whites. The rate among Hispanics was slightly higher than the rate among whites. However, the number of Hispanics living with HIV disease was small and the results should be interpreted with caution. Blacks/African Americans comprised a larger proportion of females living with HIV disease (44.1%) compared to males (12.4%). The greatest proportion of living HIV disease cases was among individuals 45 to 64 years old at the end of 2018 (47.3%).

Of the eight persons newly diagnosed with HIV disease in 2018, four were classified as stage 3 (AIDS) cases by the end of 2018 (Table 2). Among males, the majority of all new HIV disease cases diagnosed occurred among white males (50.0%). Among females, the majority of new HIV disease cases diagnosed occurred among Hispanic females (50.0%) The largest proportion of new HIV disease cases was among individuals 25 to 44 years of age (62.5%).

**Figure 6. Diagnosed and living HIV, stage 3 (AIDS), and HIV disease cases, by exposure category, Northwest HIV Care Region, 2018**



Among living HIV disease cases, the greatest proportion of cases with a known risk factor were attributed to MSM (Figure 6). The large proportion of cases with no indicated risk made trends difficult to interpret for all categories. The surveillance program examined methods to improve the identification and reporting of exposure category information.



**Table 3. New and living HIV and stage 3 (AIDS) cases and rates, by geographic area, Northwest HIV Care Region, 2018**

Geographic Area	HIV Cases						Stage 3 (AIDS) Cases					
	Diagnosed 2018*			Living			Diagnosed 2018**			Living		
	Cases	%	Rate***	Cases	%	Rate***	Cases	%	Rate***	Cases	%	Rate***
Buchanan County	4	100.0%	4.5	44	73.3%	49.5	1	25.0%	1.1	46	64.8%	51.7
Andrew County	0	0.0%	0.0	1	1.7%	5.8	0	0.0%	0.0	3	4.2%	17.3
Caldwell County	0	0.0%	0.0	1	1.7%	11.0	2	50.0%	22.1	6	8.5%	66.2
Nodaway County	0	0.0%	0.0	4	6.7%	17.6	0	0.0%	0.0	4	5.6%	17.6
Remainder of Region	0	0.0%	0.0	10	16.7%	11.7	1	25.0%	1.2	12	16.9%	14.0
<b>NORTHWEST HIV CARE REGION</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>60</b>	<b>100.0%</b>	<b>26.8</b>	<b>4</b>	<b>100.0%</b>	<b>1.8</b>	<b>71</b>	<b>100.0%</b>	<b>31.7</b>

\*HIV cases diagnosed and reported to DHSS during 2018 which remained HIV cases at the end of the year.

\*\*Does not include HIV cases diagnosed prior to 2018 that progressed to stage 3 (AIDS) in 2018.

\*\*\*Per 100,000 population based on 2017 DHSS estimates.

Note: Percentages may not total 100% due to rounding.

The greatest proportion of living HIV disease cases was diagnosed in Buchanan County (Table 3). In Buchanan County, 51.7% of living HIV disease cases progressed to stage 3 (AIDS) by the end of 2018. The rate of individuals living with HIV and stage 3 (AIDS) was also greatest in Buchanan County.

**Table 4. Newly diagnosed and living HIV and stage 3 (AIDS) cases in men who have sex with men, by selected race/ethnicity, Northwest HIV Care Region, 2018**

Race/Ethnicity	HIV Cases*				Stage 3 (AIDS) Cases			
	Newly Diagnosed		Living		Newly Diagnosed**		Living	
	Cases	%	Cases	%	Cases	%	Cases	%
White	0	--	34	89.5%	1	100.0%	23	88.5%
Black/African American	0	--	3	7.9%	0	0.0%	2	7.7%
Hispanic	0	--	1	2.6%	0	0.0%	1	3.8%
Other/Unknown	0	--	0	0.0%	0	0.0%	0	0.0%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>0</b>	<b>--</b>	<b>38</b>	<b>100.0%</b>	<b>1</b>	<b>100.0%</b>	<b>26</b>	<b>100.0%</b>

\*Remained HIV cases at the end of the year.

\*\*Does not include HIV cases diagnosed prior to 2018 that progressed to stage 3 (AIDS) in 2018.

Note: Percentages may not total 100% due to rounding.

**Table 5. Living HIV disease cases in men who have sex with men, by selected race/ethnicity and current age group, Northwest HIV Care Region, 2018**

Age Group	White		Black/African American		Hispanic		Total*	
	Cases	%**	Cases	%**	Cases	%**	Cases	%**
13-18	0	0.0%	0	0.0%	0	0.0%	0	0.0%
19-24	3	5.3%	1	20.0%	0	0.0%	4	6.3%
25-44	20	35.1%	2	40.0%	2	100.0%	24	37.5%
45-64	23	40.4%	2	40.0%	0	0.0%	25	39.1%
65+	11	19.3%	0	0.0%	0	0.0%	11	17.2%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>57</b>	<b>100.0%</b>	<b>5</b>	<b>100.0%</b>	<b>2</b>	<b>100.0%</b>	<b>64</b>	<b>100.0%</b>

\*Row totals and percentages include cases in persons whose race/ethnicity is either unknown or not listed.

\*\*Percentage of cases per age group.

Note: Percentages may not total 100% due to rounding.

**Table 6. Living HIV disease cases in men who have sex with men, by geographic area, Northwest HIV Care Region, 2018**

Geographic Area	Cases	%
Buchanan County	46	71.9%
Remaining Counties	18	28.1%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>64</b>	<b>100.0%</b>

Note: Percentages may not total 100% due to rounding.

One new HIV disease diagnoses was attributed to MSM in 2018 for the Northwest HIV Care Region (Table 4). There were 64 living HIV disease cases attributed to MSM in the Northwest HIV Care Region. Whites represented 89.5% of living HIV cases and 88.5% of living stage 3 (AIDS) cases.

The distribution of living HIV disease cases by current age varied by race/ethnicity among MSM (Table 5). Among white MSM living with HIV disease, the greatest proportion was between 45 and 64 years of age at the end of 2018. The greatest proportions of black/African American MSM living with HIV disease were 25 to 44 years of age and 45 to 64 years of age.

Buchanan County residents accounted for the largest number of living MSM in the Northwest HIV Care Region (Table 6).

**Table 7. Newly diagnosed and living HIV and stage 3 (AIDS) cases in men who have sex with men and inject drugs, by selected race/ethnicity, Northwest HIV Care Region, 2018**

Race/Ethnicity	HIV Cases*				Stage 3 (AIDS) Cases			
	Newly Diagnosed		Living		Newly Diagnosed**		Living	
	Cases	%	Cases	%	Cases	%	Cases	%
White	0	--	2	100.0%	0	--	7	100.0%
Black/African American	0	--	0	0.0%	0	--	0	0.0%
Hispanic	0	--	0	0.0%	0	--	0	0.0%
Other/Unknown	0	--	0	0.0%	0	--	0	0.0%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>0</b>	<b>--</b>	<b>2</b>	<b>100.0%</b>	<b>0</b>	<b>--</b>	<b>7</b>	<b>100.0%</b>

\*Remained HIV cases at the end of the year.  
 \*\*Does not include HIV cases diagnosed prior to 2018 that progressed to stage 3 (AIDS) in 2018.  
 Note: Percentages may not total 100% due to rounding.

**Table 8. Living HIV disease cases in men who have sex with men and inject drugs, by selected race/ethnicity and current age group, Northwest HIV Care Region, 2018**

Age Group	White		Black/African American		Hispanic		Total*	
	Cases	%**	Cases	%**	Cases	%**	Cases	%**
13-18	0	0.0%	0	--	0	--	0	0.0%
19-24	0	0.0%	0	--	0	--	0	0.0%
25-44	1	11.1%	0	--	0	--	1	11.1%
45-64	6	66.7%	0	--	0	--	6	66.7%
65+	2	22.2%	0	--	0	--	2	22.2%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>9</b>	<b>100.0%</b>	<b>0</b>	<b>--</b>	<b>0</b>	<b>--</b>	<b>9</b>	<b>100.0%</b>

\*Row totals and percentages include cases in persons whose race/ethnicity is either unknown or not listed.  
 \*\*Percentage of cases per age group.  
 Note: Percentages may not total 100% due to rounding.

**Table 9. Living HIV disease cases in men who have sex with men and inject drugs, Northwest HIV Care Region, 2018**

Geographic Area	Cases	%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>9</b>	<b>100.0%</b>

No new HIV disease diagnoses were attributed to MSM/IDU in 2018 for the Northwest HIV Care Region (Table 7). There were nine MSM/IDU living with HIV disease at the end of 2018 whose most recent diagnosis occurred in the Northwest HIV Care Region. Whites represented all of the living HIV and stage 3 (AIDS) cases.

Overall, the majority of MSM/IDU living with HIV disease (66.7%) were between 45 and 64 years of age at the end of 2018 (Table 8).

**Table 10. Newly diagnosed and living HIV and stage 3 (AIDS) cases in injection drug users, by selected race/ethnicity and sex, Northwest HIV Care Region, 2018**

Race/Ethnicity and Sex	HIV Cases*				Stage 3 (AIDS) Cases			
	Newly Diagnosed		Living		Newly Diagnosed**		Living	
	Cases	%	Cases	%	Cases	%	Cases	%
White Male	0	--	0	0.0%	0	--	2	33.3%
Black/African American Male	0	--	0	0.0%	0	--	2	33.3%
Hispanic Male	0	--	0	0.0%	0	--	0	0.0%
White Female	0	--	1	100.0%	0	--	2	33.3%
Black/African American Female	0	--	0	0.0%	0	--	0	0.0%
Hispanic Female	0	--	0	0.0%	0	--	0	0.0%
<b>NORTHWEST HIV CARE REGION TOTAL†</b>	<b>0</b>	<b>--</b>	<b>1</b>	<b>100.0%</b>	<b>0</b>	<b>--</b>	<b>6</b>	<b>100.0%</b>

\*Remained HIV cases at the end of the year.

\*\*Does not include HIV cases diagnosed prior to 2018 that progressed to stage 3 (AIDS) in 2018.

†Includes persons whose race/ethnicity is either unknown or not listed.

Note: Percentages may not total 100% due to rounding.

**Table 11. Living HIV disease cases in injection drug users, by selected race/ethnicity and current age group, Northwest HIV Care Region, 2018**

Age Group	White Males		Black/African American Males		White Females		Black/African American Females		Total*	
	Cases	%**	Cases	%**	Cases	%**	Cases	%**	Cases	%**
	13-18	0	0.0%	0	0.0%	0	0.0%	0	--	0
19-24	0	0.0%	0	0.0%	0	0.0%	0	--	0	0.0%
25-44	0	0.0%	1	50.0%	1	33.3%	0	--	2	28.6%
45-64	2	100.0%	1	50.0%	2	66.7%	0	--	5	71.4%
65+	0	0.0%	0	0.0%	0	0.0%	0	--	0	0.0%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>2</b>	<b>100.0%</b>	<b>2</b>	<b>100.0%</b>	<b>3</b>	<b>100.0%</b>	<b>0</b>	<b>--</b>	<b>7</b>	<b>100.0%</b>

\*Row totals and percentages include cases in persons whose race/ethnicity is either unknown or not listed.

\*\*Percentage of cases per age group.

Note: Percentages may not total 100% due to rounding.

**Table 12. Living HIV disease cases in injection drug users, Northwest HIV Care Region, 2018**

Geographic Area	Cases	%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>7</b>	<b>100.0%</b>

No new HIV disease diagnoses were attributed to IDU in 2018 for the Northwest HIV Care Region (Table 10). There were six living HIV disease cases attributed to IDU at the end of 2018 in the Northwest HIV Care Region. Of the living HIV disease cases, 85.7% were classified as stage 3 (AIDS) at the end of 2018. Males represented 66.7% of the living cases among IDU.

Among IDU living with HIV disease, the majority (71.4%) were 45 to 64 years of age at the end of 2018 (Table 11).

**Table 13. Newly diagnosed and living HIV and stage 3 (AIDS) cases in heterosexual contacts, by selected race/ethnicity and sex, Northwest HIV Care Region, 2018**

Race/Ethnicity and Sex	HIV Cases*				Stage 3 (AIDS) Cases			
	Newly Diagnosed		Living		Newly Diagnosed**		Living	
	Cases	%	Cases	%	Cases	%	Cases	%
White Male	0	0.0%	1	10.0%	0	--	0	0.0%
Black/African American Male	0	0.0%	0	0.0%	0	--	1	6.7%
Hispanic Male	0	0.0%	0	0.0%	0	--	0	0.0%
White Female	1	50.0%	5	50.0%	0	--	7	46.7%
Black/African American Female	0	0.0%	3	30.0%	0	--	7	46.7%
Hispanic Female	1	50.0%	1	10.0%	0	--	0	0.0%
<b>NORTHWEST HIV CARE REGION TOTAL<sup>†</sup></b>	<b>2</b>	<b>100.0%</b>	<b>10</b>	<b>100.0%</b>	<b>0</b>	<b>--</b>	<b>15</b>	<b>100.0%</b>

\*Remained HIV cases at the end of the year.

\*\*Does not include HIV cases diagnosed prior to 2018 that progressed to stage 3 (AIDS) in 2018.

†Includes persons whose race/ethnicity is either unknown or not listed.

Note: Percentages may not total 100% due to rounding.

**Table 14. Living HIV disease cases in heterosexual contacts, by selected race/ethnicity and sex and current age group, Northwest HIV Care Region, 2018**

Age Group	White Males		Black/African American Males		White Females		Black/African American Females		Total*	
	Cases	%**	Cases	%**	Cases	%**	Cases	%**	Cases	%**
	13-18	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0
19-24	0	0.0%	0	0.0%	0	0.0%	1	10.0%	1	4.0%
25-44	0	0.0%	1	100.0%	6	50.0%	3	30.0%	11	44.0%
45-64	0	0.0%	0	0.0%	5	41.7%	6	60.0%	11	44.0%
65+	1	100.0%	0	0.0%	1	8.3%	0	0.0%	2	8.0%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>1</b>	<b>100.0%</b>	<b>1</b>	<b>100.0%</b>	<b>12</b>	<b>100.0%</b>	<b>10</b>	<b>100.0%</b>	<b>25</b>	<b>100.0%</b>

\*Row totals and percentages include cases in persons whose race/ethnicity is either unknown or not listed.

\*\*Percentage of cases per age group.

Note: Percentages may not total 100% due to rounding.

**Table 15. Living HIV disease cases in heterosexual contacts, by geographic area, Northwest HIV Care Region, 2018**

Geographic Area	Total	
	Cases	%
Buchanan County	19	76.0%
Remaining Counties	6	24.0%
<b>NORTHWEST HIV CARE REGION TOTAL</b>	<b>25</b>	<b>100.0%</b>


Note: Percentages may not total 100% due to rounding.

Two new HIV disease diagnoses were attributed to heterosexual contact in 2018 for the Northwest HIV Care Region (Table 13). There were 25 living HIV disease cases attributed to heterosexual contact at the end of 2018 in the Northwest HIV Care Region. Of the living cases, 60% were classified as stage 3 (AIDS) at the end of 2018. Females represented 90% of the living HIV disease cases.


At the end of 2018, persons 25 to 44 years of age and persons 45-64 years of age comprised the largest number of heterosexual contact cases living with HIV disease in the Northwest HIV Care Region (Table 14).

At the end of 2018, Buchanan County had the largest number of heterosexual contact cases living with HIV disease in the Northwest HIV Care Region (Table 15).

**Figure 7. Reported P&S syphilis cases, by race and sex and age group at diagnosis, Northwest HIV Care Region, 2018**




**Figure 8. Reported early latent syphilis cases, by race and sex and age group at diagnosis, Northwest HIV Care Region, 2018**




The only reported P&S syphilis cases were reported among white males (7) and white females (1) in 2018 in the Northwest HIV Care Region (Figure 7). There were no cases reported among black/African American males and black/African American females. From 2017 to 2018, the number of P&S syphilis cases among white males (7) and black/African American males (0) remained the same. From 2017 to 2018, the number of P&S syphilis cases decreased among white females (2 to 1), and black/African American females (1 to 0).

The only reported early latent syphilis cases reported in 2018 were among white males (4), white females (5), and black/African American males (2) (Figure 8). From 2017 to 2018, the number of early latent cases increased among white females (2 to 5), and black/African American males (1 to 2).

**Figure 9. Reported gonorrhea cases, by race and sex and age group at diagnosis, Northwest HIV Care Region, 2018**




**Figure 10. Reported chlamydia cases, by race and sex and age group at diagnosis, Northwest HIV Care Region, 2018**




The largest numbers of gonorrhea cases were reported among white males (162), followed by white females (160) (Figure 9). Among white males and white females, the largest numbers of reported cases were between 25 to 29 years of age. Among black/African American males, the largest number of reported cases were between 25 to 29 years of age and 30 to 34 years of age. Among black/African American females, the largest numbers of reported cases were diagnosed between 20 and 24 years of age.

The largest numbers of chlamydia cases were reported among white females (490) and white males (198) (Figure 10). Among white males, white females, and black/African American males, the largest numbers of reported cases were diagnosed between 20 and 24 years of age. Among black/African American females, the largest numbers of reported cases were diagnosed between 15 and 19 years of age.

**Figure 11. Reported hepatitis B cases, by sex and age group at diagnosis, Northwest HIV Care Region, 2018**



**Figure 12. Reported hepatitis C cases, by sex and age group at diagnosis, Northwest HIV Care Region, 2018**



There were 20 reported cases of hepatitis B in the Northwest HIV Care Region during 2018 (Figure 11). Males represented 70% of reported hepatitis B cases. Among males, the largest numbers of reported cases were equal between 30 to 39 years of age and 40 to 49 years of age. The largest numbers of reported cases among females were between 30 and 39 years of age.

In 2018, there were 216 hepatitis C cases reported in the Northwest HIV Care Region (Figure 12). Of the reported hepatitis C cases, 69.0% were male. Among males, the largest numbers of reported cases were equal between 50 to 59 years of age and 60+ of age. The largest numbers of reported cases were between 50 and 59 years of age among females.



This page was intentionally left blank.