

# Missouri Weekly Influenza Surveillance Report 2017-2018 Influenza Season<sup>1</sup>

Week 3: January 14 - 20, 2018

All data are preliminary and may change as more reports are received.

## **Summary:**

- The estimated influenza activity in Missouri is Widespread<sup>2</sup>.
- During Week 3, a total of 8,260 laboratory-positive<sup>3</sup> influenza cases (5,983 influenza A, 2,193 influenza B, and 84 untyped) were reported. A season-to-date total of 53,769 laboratory-positive influenza cases (42,965 influenza A, 10,306 influenza B, and 498 untyped) have been reported in Missouri as of Week 3. The influenza type for reported season-to date cases includes 80% influenza A, 19% influenza B, and 1% untyped. Twenty-three laboratory-positive cases of influenza (17 influenza A (H3), one influenza A (H1N1), and five influenza B (Yamagata)) were reported by the Missouri State Public Health Laboratory (MSPHL) during Week 3.
- Influenza-like illness (ILI) activity was above baseline for both the Missouri Outpatient ILI Surveillance Network (ILINet) and the hospital emergency room visit chief complaint data reported through ESSENCE. The reported percentage of visits for ILI was 11.36% (Figure 5) and 6.97% (Figure 7) through ILINet and ESSENCE respectively. The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution. The percentage of respiratory specimens testing positive for influenza in Missouri laboratories reporting to the National Respiratory and Enteric Virus Surveillance System (NREVSS) decreased during Week 3 (Figure 6).
- Forty-nine influenza-associated deaths have been reported in Missouri as of Week 3.<sup>5</sup> During Week 2, 108 deaths involving Pneumonia and Influenza (P&I) were reported to the Bureau of Vital Records, resulting in a season-to-date total of 767 P&I associated deaths in Missouri.<sup>6</sup>
- Forty-eight outbreaks of influenza or ILI have been reported and five influenza or ILI-associated school closures have been reported in Missouri as of Week 3.
- Influenza activity increased in the U.S. during Week 2. National influenza surveillance information is prepared by CDC and is included in the weekly FluView report, which is available online at <a href="http://www.cdc.gov/flu/weekly/fluactivitysurv.htm">http://www.cdc.gov/flu/weekly/fluactivitysurv.htm</a>.

<sup>&</sup>lt;sup>1</sup>The 2017-2018 influenza season begins CDC Week 40 (week ending October 7, 2017) and ends CDC Week 39 (week ending September 29, 2018).

<sup>&</sup>lt;sup>2</sup>Widespread is defined as: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

<sup>&</sup>lt;sup>3</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

<sup>&</sup>lt;sup>4</sup>Influenza-like illness (ILI) is defined by ILINet as fever (temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat without a known cause other than influenza. ILI is defined by ESSENCE as Emergency Department chief complaints for Influenza or (FeverPlus and (Cough or SoreThroat) and not NonILIFevers).

<sup>&</sup>lt;sup>5</sup>All influenza-associated deaths became reportable in Missouri in 2016.

<sup>&</sup>lt;sup>6</sup>The P&I data are available one week later. The P&I data for the CDC Week provided is the most current data available.

# **Surveillance Data:**

## **Interactive Maps**

The jurisdiction-specific influenza data are provided though interactive maps available at <a href="http://arcg.is/15mPvK">http://arcg.is/15mPvK</a>. Click on the jurisdiction to view the influenza data specific to that jurisdiction.

- Reported Laboratory-positive Influenza Cases by Influenza Type by Jurisdiction, CDC Week 3
- Reported Week-specific Rate per 100,000 Population, CDC Week 3
- Reported Laboratory-positive Influenza Cases by Influenza Type by Jurisdiction, Season-to-Date
- Reported Rate per 100,000 Population, Season-to-Date

### **Data Figures**

Figure 1. Number of Laboratory-positive<sup>†</sup> Influenza Cases by Influenza Type, Missouri, CDC Week 3 (January 14 – 20, 2018)<sup>\*</sup>

Influenza Type	Week 1	Week 2	Week 3	2017-2018* Season-to-Date
Influenza A	8,555	9,703	5,983	42,965
Influenza B	1,778	2,482	2,193	10,306
Influenza Unknown Or Untyped	57	148	84	498
Total	10,390	12,333	8,260	53,769

Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv).

Figure 2. Number of Laboratory-positive<sup>†</sup> Influenza Cases and Case Rates by Age Group, Missouri, CDC Week 3 (January 14 – 20, 2018)\*<sup>‡</sup>

Age Group	Week 3 Cases	Week 3 Rate <sup>‡</sup>	2017-2018* Season-to-Date	2017-2018* Season-to-Date Rate <sup>‡</sup>
00-04	1,362	363.82	8,150	2,177.05
05-24	2,845	177.31	16,595	1,034.27
25-49	1,712	89.47	11,315	591.32
50-64	1,204	97.38	8,149	659.10
65+	1,137	119.07	9,560	1,001.13
Total	8,260	135.77	53,769	883.82

Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

<sup>\*</sup>Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

<sup>\*</sup>Incidence Rate per 100,000 population


Figure 3. Number of Laboratory-positive<sup>†</sup> Influenza Cases and Case Rates by Region, Missouri, CDC Week 3 (January 14 – 20, 2018)\*<sup>‡</sup>

Region	Week 3 Cases	Week 3 Rate <sup>‡</sup>	2017-2018* Season-to-Date	2017-2018* Season-to-Date Rate <sup>‡</sup>
Central	942	139.14	6,792	1,003.25
Eastern	2,467	108.86	19,910	878.58
Northwest	2,206	138.09	11,500	719.87
Southeast	1,053	223.24	6,751	1,431.21
Southwest	1,592	148.60	8,816	822.93
Total	8,260	135.77	53,769	883.82

Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

Figure 4. Number of Laboratory-positive<sup>†</sup> Influenza Cases by CDC Week, Missouri, 2014-2018<sup>\*</sup>




<sup>&</sup>lt;sup>†</sup>Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

\*2017-2018 season-to-date through the week ending May 19, 2018 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. Week 53, 2014-2015 flu season has 3,082 reported confirmed cases. Data Source: Missouri Health Information Surveillance System (WebSurv).

Incidence Rate per 100,000 population


Figure 5. Percentage of Outpatient Visits for Influenza-like Illness (ILI), Missouri Outpatient ILI Surveillance Network (ILINet) 2014-2018\*†



<sup>\*</sup>The ILINet Region 7 (MO, IA, KS, NE) baseline is the mean percentage of patient visits for ILI during non-influenza weeks for the previous three seasons, plus two standard deviations. A non-influenza week is defined as periods of two or more consecutive weeks in which each week accounted for less than 2% of the season's total number of specimens that tested positive for influenza.


Data Source: U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Centers for Disease Control and Prevention (CDC).

<sup>&</sup>lt;sup>†</sup>2017-2018 season-to-date through the week ending May 19, 2018 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of outpatient visits for ILI during Week 53 was 7.63.



2016-2017 season-to-date through the week ending May 19, 2018 (Week 20).


Figure 7. Percentage of Emergency Department (ED) Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, 2014-2018 Influenza Seasons\*†‡



\*The ESSENCE ILI Baseline is the mean percent of ILI visits for each week during the previous three years (2014-16) when percentage of ILI visits were less than 2% of total visits, plus two standard deviations.

Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20.

Figure 8. Percentage of Emergency Department (ED) Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by Age Group, Region and Statewide, Week 3, 2018\*




Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20. 
\*The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.

<sup>&</sup>lt;sup>†</sup>The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of visits for ILI in ESSENCE participating hospitals during Week 53 was 4.38%.


<sup>&</sup>lt;sup>‡</sup>The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.

Figure 9. Percentage of Emergency Department (ED) Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by Region and Statewide, 2017-2018 Influenza Season\*




Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20. 
\*The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.

Figure 10. Weekly Rate of Patients Hospitalized with Influenza and/or Pneumonia Syndromes in Missouri Hospitals, 2014-2018 Influenza Seasons



Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, HL7 Messaging Portal. Population data from DHSS Population MICA 2015 (http://health.mo.gov/data/mica/population.php).

Figure 11. Number of Patients Hospitalized with Influenza and/or Pneumonia Syndromes in Participating Missouri Hospitals, by Age Group, Week 3, 2017-2018 Influenza Season



Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, HL7 Messaging Portal.

#### **Additional Influenza Data Sources:**

Centers for Disease Control and Prevention: National Influenza Surveillance (FluView): http://www.cdc.gov/flu/weekly/fluactivitysurv.htm

The National Respiratory and Enteric Virus Surveillance System (NREVSS): <a href="https://www.cdc.gov/surveillance/nrevss/">https://www.cdc.gov/surveillance/nrevss/</a>

World Health Organization: International Influenza Surveillance: http://www.who.int/influenza/surveillance\_monitoring/en/