

Missouri Weekly Influenza Surveillance Report 2017-2018 Influenza Season¹

Week 9: February 25 – March 3, 2018

All data are preliminary and may change as more reports are received.

Summary:

- The estimated influenza activity in Missouri is Widespread².
- During Week 9, a total of 3,117 laboratory-positive³ influenza cases (1,304 influenza A, 1,741 influenza B, and 72 untyped) were reported. A season-to-date total of 119,071 laboratory-positive influenza cases (79,786 influenza A, 37,594 influenza B, and 1,691 untyped) have been reported in Missouri as of Week 9. The influenza type for reported season-to date cases includes 67% influenza A, 32% influenza B, and 1% untyped. Fourteen laboratory-positive cases of influenza (two influenza A (H3), two influenza A (H1N1), and 10 influenza B (Yamagata)) were reported by the Missouri State Public Health Laboratory (MSPHL) during Week 9.
- Influenza-like illness (ILI) activity was above baseline for both the Missouri Outpatient ILI Surveillance Network (ILINet) and the hospital emergency room visit chief complaint data reported through ESSENCE. The reported percentage of visits for ILI was 2.72% (Figure 5) and 3.44% (Figure 7) through ILINet and ESSENCE respectively.⁴ The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution. The percentage of respiratory specimens testing positive for influenza in Missouri laboratories reporting to the National Respiratory and Enteric Virus Surveillance System (NREVSS) decreased during Week 9 (Figure 6).
- One hundred and ninety-three influenza-associated deaths have been reported in Missouri as of Week 9.⁵ During Week 8, 91 deaths involving Pneumonia and Influenza (P&I) were reported to the Bureau of Vital Records, resulting in a season-to-date total of 1,400 P&I associated deaths in Missouri.⁶
- Sixty-six outbreaks of influenza or ILI have been reported and 15 influenza or ILI-associated school closures have been reported in Missouri as of Week 9.
- Influenza activity decreased in the U.S. during Week 8. National influenza surveillance information is prepared by CDC and is included in the weekly FluView report, which is available online at http://www.cdc.gov/flu/weekly/fluactivitysurv.htm.

⁴Influenza-like illness (ILI) is defined by ILINet as fever (temperature of 100°F [37.8°C] or greater) and a cough and/or a sore throat without a known cause other than influenza. ILI is defined by ESSENCE as Emergency Department chief complaints for Influenza or (FeverPlus and (Cough or SoreThroat) and not NonILIFevers).

⁵All influenza-associated deaths became reportable in Missouri in 2016.

⁶The P&I data are available one week later. The P&I data for the CDC Week provided is the most current data available.

¹The 2017-2018 influenza season begins CDC Week 40 (week ending October 7, 2017) and ends CDC Week 39 (week ending September 29, 2018).

²Widespread is defined as: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

³Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

Surveillance Data:

Interactive Maps

The jurisdiction-specific influenza data are provided though interactive maps available at <u>http://arcg.is/1mWHuX</u>. Click on the jurisdiction to view the influenza data specific to that jurisdiction.

- Reported Laboratory-positive Influenza Cases by Influenza Type by Jurisdiction, CDC Week 9
- Reported Week-specific Rate per 100,000 Population, CDC Week 9
- Reported Laboratory-positive Influenza Cases by Influenza Type by Jurisdiction, Season-to-Date
- Reported Rate per 100,000 Population, Season-to-Date

Data Figures

Figure 1.	Number of Laboratory-positive [†] Influenza Cases by Influenza Type,
	Missouri, CDC Week 9 (February 25 – March 3, 2018) [*]

Influenza Type	Week 7	Week 8	Week 9	2017-2018* Season-to-Date
Influenza A	5,189	2,765	1,304	79,786
Influenza B	5,423	3,773	1,741	37,594
Influenza Unknown Or Untyped	169	118	72	1,691
Total	10,781	6,656	3,117	119,071

^TLaboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture. ^{*}Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv).

Figure 2. Number of Laboratory-positive [†] Influenza Cases and Case Rates by Age Group,
Missouri, CDC Week 9 (February 25 – March 3, 2018) ^{*‡}

Age Group	Week 9 Cases	Week 9 Rate [‡]	2017-2018* Season-to-Date	2017-2018* Season-to-Date Rate [‡]
00-04	514	137.30	18,502	4,942.30
05-24	1,009	62.89	42,579	2,653.70
25-49	621	32.45	23,651	1,236.01
50-64	445	35.99	16,823	1,360.67
65+	528	55.29	17,516	1,834.29
Total	3,117	51.24	119,071	1,957.22

Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture. *Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

^{*}Incidence Rate per 100,000 population


Figure 3. Number of Laboratory-positive[†] Influenza Cases and Case Rates by Region, Missouri, CDC Week 9 (February 25 – March 3, 2018)^{*†}

Region	Week 9 Cases	Week 9 Rate [‡]	2017-2018* Season-to-Date	2017-2018* Season-to-Date Rate [‡]
Central	558	82.42	16,205	2,393.66
Eastern	938	41.39	36,270	1,600.51
Northwest	814	50.95	33,333	2,086.55
Southeast	420	89.04	13,803	2,926.23
Southwest	387	36.12	19,460	1,816.48
Total	3,117	51.24	119,071	1,957.22

[†]Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture. ^{*}Influenza season begins week ending October 7, 2017 (CDC Week 40) Data Source: Missouri Health Information Surveillance System (WebSurv)

[‡]Incidence Rate per 100,000 population





[†]Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RT-PCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

^{*}2017-2018 season-to-date through the week ending May 19, 2018 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. Week 53, 2014-2015 flu season has 3,082 reported confirmed cases. Data Source: Missouri Health Information Surveillance System (WebSurv).




Figure 5. Percentage of Outpatient Visits for Influenza-like Illness (ILI), Missouri Outpatient ILI Surveillance Network (ILINet) 2014-2018^{*†}

^{*}The ILINet Region 7 (MO, IA, KS, NE) baseline is the mean percentage of patient visits for ILI during non-influenza weeks for the previous three seasons, plus two standard deviations. A non-influenza week is defined as periods of two or more consecutive weeks in which each week accounted for less than 2% of the season's total number of specimens that tested positive for influenza.

Data Source: U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Centers for Disease Control and Prevention (CDC).

¹2017-2018 season-to-date through the week ending May 19, 2018 (Week 20). The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of outpatient visits for ILI during Week 53 was 7.63.






Figure 6. Season-to-Date PCR (+) Tests for Influenza in Missouri

Data Source: National Respiratory and Enteric Virus Surveillance System (NREVSS), Centers for Disease Control and Prevention (CDC). 2016-2017 season-to-date through the week ending May 19, 2018 (Week 20).





^{*}The ESSENCE ILI Baseline is the mean percent of ILI visits for each week during the previous three years (2014-16) when percentage of ILI visits were less than 2% of total visits, plus two standard deviations.

Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20. [†] The 2014-2015 season had 53 weeks rather than the usual 52. The percentage of visits for ILI in ESSENCE participating hospitals during Week 53 was 4.38%.

^{*} The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.




Figure 8. Percentage of Emergency Department (ED) Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by Age Group, Region and Statewide, Week 9, 2018^{*}

Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20. "The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.






Figure 9. Percentage of Emergency Department (ED) Visits for Influenza-like Illness (ILI) in ESSENCE Participating Hospitals, by Region and Statewide, 2017-2018 Influenza Season*


Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, ESSENCE version 1.20. ^{*}The ILI data from a small number of sites located in the Northwest Region of the state is temporarily unavailable in ESSENCE. Therefore, the ILI data for the Northwest Region should be interpreted with caution.

Figure 10. Weekly Rate of Patients Hospitalized with Influenza and/or Pneumonia Syndromes in Missouri Hospitals, 2014-2018 Influenza Seasons



Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, HL7 Messaging Portal. Population data from DHSS Population MICA 2015 (http://health.mo.gov/data/mica/population.php).





Data Source: Missouri Department of Health and Senior Services (DHSS), Bureau of Reportable Disease Informatics, HL7 Messaging Portal.

Additional Influenza Data Sources:

Centers for Disease Control and Prevention: National Influenza Surveillance (FluView): http://www.cdc.gov/flu/weekly/fluactivitysurv.htm

The National Respiratory and Enteric Virus Surveillance System (NREVSS): https://www.cdc.gov/surveillance/nrevss/

World Health Organization: International Influenza Surveillance: http://www.who.int/influenza/surveillance_monitoring/en/