MISSOURI MONTHLY TUBERCULOSIS DISEASE REPORT

Reported Number of Tuberculosis Disease Cases for MMWR Weeks 1-34 2016 versus 2017

| Geographic Area | Number of Counted Cases: 2016 | Number of Counted Cases: 2017 | Difference in Number of Counted Cases: 2016 to 2017 | Percent of 2017 Cases that were Preventable |
|--------------------------------------|-------------------------------------|-------------------------------------|--|---|
| | | | | |
| St. Louis City | 6 | 11 | 5 | 27% |
| St. Louis County | 16 | 9 | -7 | 0% |
| Kansas City | 9 | 7 | -2 | 14% |
| Springfield/Greene | 1 | 2 | 1 | 0% |
| Region(excluding metropolitan areas) | | | | |
| Northwestern | 14 | 2 | -12 | 0% |
| Central | 8 | 1 | -7 | 0% |
| Southeastern | 6 | 5 | -1 | 0% |
| Southwestern | 4 | 4 | 0 | 0% |
| Eastern | 4 | 66 | 2 | 17% |
| Institutionalized Patients* | 4 | 1 | -3 | 0% |
| TOTAL** | 72 | 48 | -24 | 10% |

^{*}Institutionalized patients are those individuals who are incarcerated in federal or state prisons. Provisional Data from the WebSurv System Subject to Change

^{**} The total percent of preventable cases is derived by dividing the total number (not percentage) of preventable cases by the total number of 2017 counted cases.